

Particle-hole condensates of higher angular momentum in hexagonal systems

Akash V. Maharaj¹, Ronny Thomale^{1,2}, and S. Raghu^{1,3}

¹*Department of Physics, Stanford University, Stanford, California 94305, USA*

²*Institut de théorie des phénomènes physiques, École Polytechnique Fédérale de Lausanne (EPFL), CH-1015 Lausanne and*

³*SLAC National Accelerator Laboratory, Menlo Park, CA 94025, USA*

(Dated: March 13, 2013)

Hexagonal lattice systems (*e.g.* triangular, honeycomb, kagome) possess a multidimensional irreducible representation corresponding to $d_{x^2-y^2}$ and d_{xy} symmetry. Consequently, various phases that break time-reversal and spin rotation symmetries can occur. We show that hexagonal lattice systems with extended repulsive interactions can exhibit instabilities in the particle-hole channel to phases with $d_{x^2-y^2} + id_{xy}$ symmetry. When lattice translational symmetry is preserved, the phase corresponds to nematic order in the spin-channel with broken time-reversal symmetry, known as the β phase. On the other hand, lattice translation symmetry can also be broken, resulting in various $d_{x^2-y^2} + id_{xy}$ density wave orders. In the weak-coupling limit, when the Fermi surfaces lie close to a van Hove singularity, instabilities of both types are obtained in a controlled fashion.

Introduction.- Correlated electron materials often exhibit a tendency towards forming multiple, competing phases including superconductivity, density waves, and phases with orientational order. In the weak-coupling limit, the set of all possible broken symmetries is determined by symmetry alone; each is labeled by an allowed irreducible representation (irrep) of the symmetric normal state. When symmetry allows for higher dimensional irreps, phases that spontaneously break time-reversal and/or spin rotation symmetry can occur. Examples include a $p_x + ip_y$ superconductor in a tetragonal system, or the B-phase of Helium-3 in a cubic crystal. The former breaks time-reversal, whereas the latter represents a phase with broken spin rotation symmetry, *i.e.* an example of dynamically generated spin-orbit coupling.

Here, we study systems with hexagonal symmetry. There, $d_{x^2-y^2}$ and d_{xy} irreps are degenerate, forming a two-component irrep [1]. This allows for the possibility of $d + id$ superconductivity, which spontaneously breaks parity and time-reversal symmetry [2–6]. However, this is not the only possibility: hexagonal systems can also naturally allow for *particle-hole* condensates with $d + id$ symmetry. When such a system preserves lattice translation symmetry, the state is known as a β phase [7]: an electron nematic state in the spin channel, which spontaneously breaks time-reversal and spin-rotation symmetry. In this phase, spin orbit coupling is generated dynamically [8]. When the particle-hole condensate breaks translation symmetry, it is known as a $d + id$ density wave phase and can occur either in the spin singlet or triplet channel [9].

In this paper, we demonstrate the existence of both the β phase, and $d + id$ density wave ground state on a triangular lattice. We show here that both phases are stabilized in the weak-coupling limit when the Fermi surface is tuned to cross a van Hove singularity. In such a limit, there is an instability to each of these phases allowing the use of an unrestricted Hartree-Fock treatment. Although this fine tuning is required to access the instabilities in

a controlled fashion, it is likely that such phases survive away from the van Hove filling in the experimentally relevant intermediate coupling regime. We also study the transition between the two phases as the degree of Fermi surface nesting is altered. Since both phases represent particle-hole condensates of higher relative angular momentum, they require longer range interactions in order to be stabilized over their zero angular momentum counterparts - namely ferromagnetism and spin density wave phases, respectively [10]. Since the considerations below are based solely on symmetry, the results are also appropriate on other hexagonal lattices such as the honeycomb and kagome, as has been confirmed by recent studies [5, 11–16].

Extended Hubbard model.- We consider spinful fermions hopping on a triangular lattice, with nearest and next nearest neighbor (nnn) hopping strengths, t_1 and t_2 , and an onsite and nearest neighbor density-density interaction. The Hamiltonian is

$$H = \sum_{\mathbf{k}, \sigma} \varepsilon_{\mathbf{k}} c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma} + U \sum_i n_{i\uparrow} n_{i\downarrow} + V \sum_{\langle i,j \rangle} \sum_{\sigma, \sigma'} n_{i\sigma} n_{j\sigma'}, \quad (1)$$

where $n_{i\sigma} = c_{i\sigma}^\dagger c_{i\sigma}$ is the density operator for electrons of spin σ , at site i . The dispersion is $\varepsilon_{\mathbf{k}} = -2t_1 [\cos k_x + 2 \cos(k_x/2) \cos(\sqrt{3}k_y/2)] - 2t_2 [\cos \sqrt{3}k_y + 2 \cos(3k_x/2) \cos(\sqrt{3}k_y/2)]$ and there is a logarithmic divergence in the density of states at the van Hove filling of $\mu^* = 2(t_1 + t_2)$. Fourier transforming the interaction yields

$$H_{\text{int}} = -U \sum_{\mathbf{k}, \mathbf{k}', \mathbf{q}} c_{\mathbf{k}\downarrow}^\dagger c_{\mathbf{k}+\mathbf{q}\uparrow} c_{\mathbf{k}'+\mathbf{q}\uparrow}^\dagger c_{\mathbf{k}'\downarrow} - V \sum_{\mathbf{k}, \mathbf{k}', \mathbf{q}} \sum_{\sigma, \sigma'} f_{\mathbf{k}, \mathbf{k}'} c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}+\mathbf{q}\sigma} c_{\mathbf{k}'+\mathbf{q}\sigma}^\dagger c_{\mathbf{k}'\sigma'}. \quad (2)$$

From symmetry considerations, it follows that $f_{\mathbf{k}, \mathbf{k}'}$, which are lattice analogs of Landau parameters, can be written as a sum of separable interactions in each of

the distinct irreps of the normal state point group. For hexagonal systems with a D_{6h} point group, this sum can be written as

$$f_{\mathbf{k},\mathbf{k}'} = \frac{1}{3}d_{\mathbf{k}}^{(A_{1g})}d_{\mathbf{k}'}^{(A_{1g})} + \frac{2}{3}\mathbf{d}_{\mathbf{k}}^{(E_{2g})} \cdot \mathbf{d}_{\mathbf{k}'}^{(E_{2g})} + \frac{1}{3}d_{\mathbf{k}}^{(B_{2u})}d_{\mathbf{k}'}^{(B_{2u})} + \frac{2}{3}\mathbf{d}_{\mathbf{k}}^{(E_{1u})} \cdot \mathbf{d}_{\mathbf{k}'}^{(E_{1u})}, \quad (3)$$

where $d_{\mathbf{k}}^{(A_{1g})}$ is the explicit form of the A_{1g} or extended s -wave representation etc. The other irreducible representations present are f -wave, and the two doubly degenerate representations relate to d -wave and p -wave. For the E_{2g} correlations which we are interested in, the explicit $\{d_{x^2-y^2}, d_{xy}\}$ form factors are $\{\cos k_x - \cos \frac{k_x}{2} \cos \frac{\sqrt{3}k_y}{2}, \sqrt{3} \sin \frac{k_x}{2} \sin \frac{\sqrt{3}k_y}{2}\}$.

Fermi surface instabilities.- For purely repulsive interactions, there is no superconducting instability to first-order in the interactions - *i.e.* at mean field level - since there are no negative eigenvalues of the interaction matrix. We therefore examine instabilities in the particle-hole channel towards the formation of phases with order parameters $\Delta_{\mathbf{q};\sigma,\sigma'}^{(\eta)} = \sum_{\mathbf{k}} V^{(\eta)} d_{\mathbf{k}}^{(\eta)} \langle c_{\mathbf{k}+\mathbf{q}\sigma}^\dagger c_{\mathbf{k}\sigma'} \rangle$, where η labels the irrep. Having decomposed the interaction into separate irreps, we see that in the Random Phase Approximation (RPA), the effective interaction will not mix different irreps. The effective interaction in RPA, $\Gamma_{\mathbf{k},\mathbf{k}'}$, will have the form

$$\Gamma_{\mathbf{k},\mathbf{k}'} = \sum_{\eta} \frac{V^{(\eta)}}{1 - V^{(\eta)}\chi^{(\eta)}(\mathbf{q})} d_{\mathbf{k}}^{(\eta)} d_{\mathbf{k}'}^{(\eta)}, \quad (4)$$

where $\chi^{(\eta)}(\mathbf{q})$ are the non-interacting susceptibilities, and the sum is over *distinct* irreps η as in Eq. 3. The free susceptibility for each irrep diverges logarithmically at the van Hove filling, so the dominant instability is given by the one with the largest pre-factor.

Mean Field Analysis.- When the Fermi surface is perfectly nested, hexagonal symmetry dictates that there are three nesting vectors $\mathbf{Q}_{1,2,3}$ as defined in Fig. 1a. Thus, $\chi(\mathbf{q})$ will be most divergent for any of these three vectors and this corresponds to charge or spin density wave order. However, as shown in Fig. 1, finite second neighbor hopping destroys this perfect nesting and we may expect $\mathbf{q} = 0$ orders to be favored beyond some critical value of t_2 . We will see that nematic instabilities are dominant for large enough V and t_2 . While such instabilities can occur in the charge channel, there is the more exotic possibility of realizing a $d + id$ nematic phase in the spin channel. We must invoke the spin degrees of freedom since the Hamiltonian must remain Hermitian, and we gain condensation energy utilizing both irreps. This so called β phase is in fact the most energetically preferable instability, and the corresponding mean field




FIG. 1. (a). The Fermi surface of the un-ordered state at the van Hove filling as the second neighbor hopping t_2 is tuned from negative (blue), to zero (black) where it is perfectly nested along three vectors $\mathbf{Q}_{1,2} = \pi(1, \pm\sqrt{3})$ and $\mathbf{Q}_3 = 2\pi(1, 0)$, and finally to positive (pink). (b) The Fermi surface splitting from the original black surface, into red and blue contours upon entering the $d + id$ Pomeranchuk phase; the van Hove points are removed, and the spin winds twice around the Brillouin zone. The spin orientation is opposite on each Fermi surface, and the inner circle shows winding on one of these.

Hamiltonian is

$$H = \sum_{\mathbf{k}} \Psi_{\mathbf{k}}^\dagger \left[\epsilon_{\mathbf{k}} \mathbb{I}_2 - \Delta \left(d_{\mathbf{k}}^{(1)} \tau^1 + d_{\mathbf{k}}^{(2)} \tau^2 \right) \right] \Psi_{\mathbf{k}} + \frac{|\Delta|^2}{(2V/3)}, \quad (5)$$

where, $\Psi_{\mathbf{k}}^\dagger = (c_{\mathbf{k},\uparrow}^\dagger, c_{\mathbf{k},\downarrow}^\dagger)$, and τ^i are the usual Pauli matrices in the spin space. Δ is the magnitude of the order parameter written above, while $d_{\mathbf{k}}^{(i)}$ are the d -wave form factors. By diagonalizing this matrix, we see that the Fermi surface splits into two parts, with the spin winding twice in momentum space (see Fig. 1b). For the perfectly nested Fermi surface, the $d + id$ density wave state may occur in either the singlet or triplet channels, with a mean field Hamiltonian of similar form to Eq. 5, but with $\Psi_{\mathbf{k}}^\dagger = (c_{\mathbf{k}}^\dagger, c_{\mathbf{k}+\mathbf{Q}}^\dagger)$, where $\mathbf{Q}$ is the commensurate ordering vector, and the Pauli matrices no longer acting in spin space.

The static susceptibility for a particular irrep η in the non interacting system is given by the expression

$$\chi^{(\eta)}(\mathbf{q}) = - \int \frac{d^2k}{(2\pi)^2} \frac{f(\epsilon_{\mathbf{k}+\mathbf{q}}) - f(\epsilon_{\mathbf{k}})}{\epsilon_{\mathbf{k}+\mathbf{q}} - \epsilon_{\mathbf{k}}} |d_{\mathbf{k}}^{(\eta)}|^2, \quad (6)$$

where $f(\epsilon_{\mathbf{k}})$ is the usual fermi function, and the integral runs over the first Brillouin zone. Since the zero temperature integral is dominated by the saddle points in the dispersion we can obtain analytic expressions for the Pomeranchuk and density wave susceptibilities. We then explain how these connect to a numerical evaluation of Eq. 6 at finite temperature.

For $\mathbf{q} = 0$ orders, the expression for the susceptibility becomes

$$\chi^{(\eta)}(\mathbf{q} \rightarrow 0) = \int \frac{d^2k}{(2\pi)^2} |d_{\mathbf{k}}^{(\eta)}|^2 \delta(\epsilon_{\mathbf{k}} - \mu). \quad (7)$$




FIG. 2. The coefficients of the logarithmically diverging susceptibilities at zero temperature from Eq. 8, 9, 11 and 12. This shows crossover from density wave ($\mathbf{q} = \mathbf{Q}$) to Pomeranchuk $\mathbf{q} = 0$ orders as t_2 is varied.

Following the method of Ref.[17], we can expand in polar coordinates around a single one of the six equivalent van Hove points. Choosing the point $(0, -2\pi/\sqrt{3})$, we set $k_x = r \cos \theta$ and $k_y = -2\pi/\sqrt{3} + r \sin \theta$, and so, the $\mathbf{k}$ integral can be transformed into integrals over θ and r . For the density of states (or ferromagnetic susceptibility), upon carrying out appropriate transformations on the delta functions of Eq. 7, we obtain

$$\rho(0) \sim \frac{1}{2\pi^2} \sqrt{\frac{3}{(t_1 - 9t_2)(t_1 - t_2)}} \log\left(\frac{1}{x}\right), \quad (8)$$

where $x = \mu^* - \mu$. This shows the correct logarithmic divergence at the van Hove filling of $\mu = \mu^*$. Note the van Hove points disappear at $t_2 \geq t_1/9$ when the M point is no longer a saddle, so we restrict our analysis to the range of parameter range $t_2 < t_1/9$.

On the other hand, for the d -wave Pomeranchuk susceptibility, the six van Hove points are no longer equivalent. When summed over the six saddle points, the extra factor in the integrand gives $|d_{\mathbf{k}}^{(E_{2g})}|^2 \approx 12 + \mathcal{O}(k^2)$. Thus, in the limit $\mu \rightarrow \mu^*$, the analytic expression for d -wave susceptibilities is

$$\chi^{(E_{2g})}(\mathbf{q} \rightarrow 0) \sim 2\rho(0). \quad (9)$$

Note that this is the same for both the $d_{x^2-y^2}$ and d_{xy} form factors, reflecting their symmetry protected degeneracy. For commensurate density wave order, the susceptibility integral can be rewritten as

$$\chi^{(\eta)}(\mathbf{q}) = \int \frac{d^2k}{(2\pi)^2} \frac{f(\varepsilon_{\mathbf{k}})}{\varepsilon_{\mathbf{k}+\mathbf{q}} - \varepsilon_{\mathbf{k}}} \left(|d_{\mathbf{k}}^{(\eta)}|^2 + |d_{\mathbf{k}+\mathbf{q}}^{(\eta)}|^2 \right). \quad (10)$$

For the uniform density wave at any of the nesting vectors $\mathbf{Q}$, a similar expansion in polar coordinates around each van Hove point with a cutoff to ensure $\varepsilon_{\mathbf{k}} < 0$ as dictated

by the Fermi function gives the result

$$\chi^{(0)}(\mathbf{Q}) \sim \frac{\sqrt{3}}{4\pi^2(t_1 - 3t_2)} \log\left[\frac{\tan(\theta_0 + \frac{\pi}{12})}{\tan(\frac{\pi}{12})}\right] \log\left(\frac{1}{x}\right), \quad (11)$$

where $2\theta_0$ is the angle between arcs of the Fermi surface at the M point, given by $2 \cos 2\theta_0 = (t_1 + 3t_2)/(t_1 - 3t_2)$. This also has the requisite logarithmic divergence at the van Hove filling, but with a different coefficient to the density of states.

Note that the d_{xy} and $d_{x^2-y^2}$ DW susceptibilities are no longer degenerate since the point group symmetry is broken by a finite ordering vector. However, they do transform as a vector under the point group symmetry. The constant magnitude of this vector implies that the $d + id$ DW susceptibility is degenerate at all nesting vectors $\mathbf{Q}$, and explicit evaluation gives

$$\chi^{(d+id)}(\mathbf{Q}) = 2\chi^{(0)}(\mathbf{Q}), \quad (12)$$

which represents the same relation between d -wave and s -wave susceptibilities as for $\mathbf{q} = 0$ orders.

Since all of these susceptibilities diverge logarithmically, the dominant instability will be determined by both the interaction strength for a given irrep ($V^{(\eta)}$), and the coefficient of the diverging susceptibility. These coefficients are plotted in Fig. 2, showing that at $t_2 \geq 0.04t_1$, the $\mathbf{q} = 0$ orders become dominant. These analytic results at zero temperature provide valuable insight into the finite temperature values of the susceptibility. At finite temperatures the logarithmic divergences are tamed, but susceptibilities are still peaked strongly at the van Hove filling. Numerical evaluation of these finite T susceptibilities shows that the maximum values of χ are proportional to coefficients shown in Fig. 2. We can therefore determine the phase diagram by using a Stoner criterion for each phase. The result is a rich phase diagram in the plane of V vs. t_2 , with $l = 0$ orders dominant at small values of V , while higher angular momentum $l = 2$ particle hole condensates are favored when longer range interactions are larger. The transition between these phases takes place at a critical nearest neighbor V given by $V^{E_{2g}} \chi^{E_{2g}}(\mathbf{q}) \equiv \frac{2}{3} V^* \chi^{E_{2g}}(\mathbf{q}) > U \chi^0(\mathbf{q})$, which is around $V^* \sim 0.8U$. Numerical minimization of the mean field free energy show that the transitions between these phases are first order at zero temperature, but the transition into each phase is continuous as a function of temperature.

Landau-Ginzburg theory.- Having determined that the nematic and dDW instabilities are dominant for large V , we now ask what combination of the $d_{x^2-y^2}$ and d_{xy} order parameters is preferable. Our numerical calculations confirm that these transitions are continuous as a function of temperature, so we can safely truncate a Landau-Ginzburg free energy expansion at low order. The presence of a two component order parameter along with the




FIG. 3. Phase diagram of the model in Eq. 1 as a function of nearest neighbor interaction V and second neighbor hopping t_2 . At low values of V , the uniform or zero angular momentum particle hole condensates dominate, while the time reversal symmetry breaking $l = 2$ condensates dominate above a critical V/U . All phase boundaries are first order lines at zero temperature.

C_{6v} symmetry, dictates that the expansion for the $\mathbf{q} = 0$ Pomeranchuk order parameters must be of the form

$$\mathcal{F} = a(T - T_c)(|\Delta_1|^2 + |\Delta_2|^2) + b_1(|\Delta_1|^2 + |\Delta_2|^2)^2 + b_2(\Delta_1^* \Delta_2 - \Delta_1 \Delta_2^*)^2, \quad (13)$$

up to fourth order [1], where Δ_1 and Δ_2 are the $d_{x^2-y^2}$ and d_{xy} components of the order parameter. The sign of the coefficient b_2 determines which combination of the order parameters is present; for $b_2 > 0$, the sum $\Delta_1 \pm i\Delta_2$ minimizes the free energy, whereas for $b_2 < 0$ either a d_{xy} or $d_{x^2-y^2}$ order results. These coefficients can be determined microscopically [18], and in a manner exactly analogous with superconductivity [4], we get $b_1 = \frac{1}{2}K$ and $b_2 = \frac{1}{12}K$ where $K > 0$ is a trace over four Green functions. Thus, the $d + id$ Pomeranchuk state is favored at the transition, and numerical minimization of the mean field free energy suggests that this state persists deep below T_c since it gains maximum condensation energy. Similar considerations favor the $d + id$ density wave state.

Discussion.- We have shown on analytical footing how higher angular momentum particle hole condensates can be realized from models with extended interactions on hexagonally symmetric lattices. The symmetry protected degeneracy of the d -wave representations means that the $d + id$ time reversal symmetry breaking phases are preferred, in a manner that is analogous to superconductivity.

The full phase diagram in the plane V vs. t_2 is shown in Fig. 3. This provides a unified view of the effects of longer range interactions as bandstructure effects are tuned. When the Fermi surface is approximately perfectly nested, commensurate density wave order is dominant. For low values of V , a spin density wave forms

at any of the three nesting vectors or some combination of them [15]. Meanwhile, higher angular momentum $d + id$ density waves are favored by longer ranged interactions. This can be either a singlet or triplet density wave, with the triplet states accompanied by a Goldstone boson due to the broken spin rotation symmetry. Furthermore, there can be an analog of the time reversal symmetry preserving quantum spin Hall state, with opposite spins winding with different chirality ($d + id$ for up and $d - id$ for down spins respectively)[19].

However, as the nesting is destroyed by finite t_2 , the $\mathbf{q} = 0$ orders are favored, with an analogous crossover from $l = 0$ ferromagnetism to an $l = 2$ chiral nematic phase as the interaction strength is increased. The latter is the β phase - a $d + id$ Pomeranchuk instability which occurs in the spin channel. This is a metal with dynamically driven spin orbit coupling, i.e. one that spontaneously breaks time reversal and $SU(2)$ spin rotation symmetries. We note that these Pomeranchuk phases have been observed as same sub-lattice instabilities in FRG studies of Kagome lattices, which would suggest that these infinitesimal interaction results may persist at stronger coupling [13].

While we have considered here the particular case of hexagonal systems, it is likely that non-trivial particle hole condensates can occur in cubic lattices which allow for three dimensional t_{2g} irreps. It is likely, therefore, that the β phase, which makes use of each of the irreps in the spin channel will be the favored ground state at a Pomeranchuk instability. We shall present the analysis of three dimensional systems elsewhere.

We acknowledge insightful discussions with Andrey Chubukov and Shou-Cheng Zhang. RT thanks M. Kiesel and C. Platt for collaborations on related topics. This work was supported in part by the U.S. Department of Energy under Contract No. AC02-76SF00515 (AM,SR), and the Alfred P. Sloan Foundation (SR).

-
- [1] M. Sigrist and K. Ueda, Rev. Mod. Phys. **63**, 239 (1991).
 - [2] C. Honerkamp, Phys. Rev. Lett. **100**, 146404 (2008).
 - [3] S. Raghu, S. A. Kivelson, and D. J. Scalapino, Phys. Rev. B **81**, 224505 (2010).
 - [4] R. Nandkishore, L. S. Levitov, and A. V. Chubukov, Nature Physics **8**, 158 (2012).
 - [5] M. L. Kiesel, C. Platt, W. Hanke, D. A. Abanin, and R. Thomale, Phys. Rev. B **86**, 020507 (2012).
 - [6] M. Kiesel, C. Platt, W. Hanke, and R. Thomale, arXiv:1301.5662.
 - [7] C. Wu, K. Sun, E. Fradkin, and S.-C. Zhang, Phys. Rev. B **75**, 115103 (2007).
 - [8] C. Wu and S.-C. Zhang, Physical review letters **93**, 36403 (2004).
 - [9] C. Nayak, Physical Review B **62**, 4880 (2000).
 - [10] H. J. Schulz, Phys. Rev. B **39**, 2940 (1989).
 - [11] W. Wang, Y. Xiang, Q. Wang, F. Wang, F. Yang, and

- D. Lee, Physical Review B **85**, 035414 (2012).
- [12] M. L. Kiesel and R. Thomale, Phys. Rev. B **86**, 121105 (2012).
- [13] M. Kiesel, C. Platt, and R. Thomale, arXiv:1209.3398, to appear in Phys. Rev. Lett.
- [14] O. Tieleman, O. Dutta, M. Lewenstein, and A. Eckardt, ArXiv e-prints (2012).
- [15] R. Nandkishore, G.-W. Chern, and A. V. Chubukov, Phys. Rev. Lett. **108**, 227204 (2012).
- [16] R. Nandkishore and A. V. Chubukov, Phys. Rev. B **86**, 115426 (2012).
- [17] H. Q. Lin and J. E. Hirsch, Phys. Rev. B **35**, 3359 (1987).
- [18] R. Fernandes and J. Schmalian, Physical Review B **82**, 014521 (2010).
- [19] C.-H. Hsu, S. Raghu, and S. Chakravarty, Phys. Rev. B **84**, 155111 (2011).